Modeling in Medical Decision Analysis

Jeremy D. Goldhaber-Fiebert, PhD

Presented February 24, 2016

HEALTH POLICY STANFORD

CENTER FOR HEALTH POLICY/
CENTER FOR PRIMARY CARE AND OUTCOMES RESEARCH

Agenda

- Decision analysis
- Cost-effectiveness analysis
- Decision trees
- Sensitivity analysis
- Markov models
- Microsimulations

POLL SLIDE

- **Have you had a course in (*select all that apply*):**
  1. Medicine
  2. Epidemiology
  3. Probability and/or Statistics
  4. Computer Programming
  5. Decision Science/Economic Evaluation
  6. Microeconomics

WHAT IS A DECISION ANALYSIS?

What is a decision analysis?

- A quantitative method for evaluating decisions between multiple alternatives in situations of uncertainty

What is a decision analysis?

- A quantitative method for evaluating decisions between multiple alternatives in situations of uncertainty

Decisions between multiple alternatives:

- Allocate resources to one alternative (and not the others)
- There is no decision without alternatives => making a choice

What is a decision analysis?

- A quantitative method for evaluating decisions between multiple alternatives in situations of uncertainty

Quantitative method for evaluating decisions:

- Gather information
- Assess the consequences of each alternative
- Clarify the dynamics and trade-offs involved in selecting each
- Select an action to take that gives us the best expected outcome

We generally employ models to do this

The steps of a decision analysis

1. Enumerate all relevant alternatives
2. Identify important outcomes
3. Determine relevant uncertain factors
4. Encode probabilities for uncertain factors
5. Specify the value of each outcome
6. Combine these elements to analyze the decision

Decision trees and related models important for this

What is a decision analysis called when its important outcomes include costs?

1. Enumerate all relevant alternatives
2. *Identify important outcomes*
3. Determine relevant uncertain factors
4. Encode probabilities for uncertain factors
5. Specify the value of each outcome
6. Combine these elements to analyze the decision

Cost-effectiveness analysis a type of decision analysis that includes costs as one of its outcomes

WHAT IS A COST-EFFECTIVENESS ANALYSIS?

What is a cost-effectiveness analysis?

- In the context of health and medicine, a *cost-effectiveness analysis (CEA)* is a method for evaluating tradeoffs between health benefits and costs resulting from alternative courses of action

- **CEA supports decision makers; it is not a complete resource allocation procedure**

Cost-Effectiveness Ratio (CER): How to compare two strategies in CEA

- **Numerator**: Difference between costs of the intervention (strategy) and costs of the
  tudy
  erence b
  eness) o
  and heal comes of the al tive

Incremental resources required by the intervention

Incremental health effects gained with the intervention

$$CER = \frac{C_i - C_{alt}}{E_i - E_{alt}}$$

Models for decision analysis and CEAs

- **Decision model:** a *schematic* representation of all of the clinically and policy relevant features of the decision problem
  - Includes the following in its structure:
    - Decision alternatives
    - Clinical and policy-relevant outcomes
    - Sequences of events
  - Enables us to integrate knowledge about the decision problem from many sources (i.e., probabilities, values)
  - Computes expected outcomes (i.e., averaging across uncertainties) for each decision alternative

Building decision-analytic model

1. Define the model's structure
2. Assign probabilities to all chance events in the structure
3. Assign values (i.e., utilities) to all outcomes encoded in the structure
4. Evaluate the expected utility of each decision alternative
5. Perform sensitivity analyses

Simple enough to be understood; complex enough to capture problem's elements convincingly (assumptions)

“All models are wrong; but some models are useful”

-- George Box and Norman Draper, 1987

Building decision-analytic models

1. *Define the model's structure*
2. Assign probabilities to all chance events in the structure
3. Assign values (i.e., utilities) to all outcomes encoded in the structure
4. Evaluate the expected utility of each decision alternative
5. Perform sensitivity analyses

WHAT ARE THE ELEMENTS OF A DECISION TREE’S STRUCTURE?

Decision node

A place in the decision tree at which there is a choice between several alternatives



The example shows a choice between 2 alternatives, but a decision node can accommodate a choice between more alternatives … provided alternatives are **mutually exclusive**.

Chance node

A place in the decision tree at which chance determines the outcome based on probability



The example shows only 2 outcomes, but a chance node can accommodate more outcomes … provided they are **<u>mutually exclusive AND collectively exhaustive</u>**.

What do <u>mutually exclusive</u> and <u>collectively exhaustive</u> mean?

- **<u>Mutually exclusive</u>**
  - Only one alternative can be chosen
  - Only one event can occur
- **<u>Collectively exhaustive</u>**
  - At least one event must occur
  - One of the possibilities must happen
  - Taken together, the possibilities make up the entire range of outcomes

Terminal node

Final outcome associated with each pathway of choices and chances



Final outcomes must be valued in relevant terms (cases of disease, Life years, Quality-adjusted life years, costs) so that they can be used for comparisons

Summary

- **Decision nodes**: enumerate a choice between alternatives for the decision maker
- **Chance nodes**: enumerate possible events determined by chance/probability
- **Terminal nodes**: describe outcomes associated with a given pathway (of choices and chances)

The entire structure of the decision tree can be described with only these elements

Example: decision tree

- Patient presents with symptoms
- Likely serious disease; unknown w/o treatment
- Two treatment alternative:
  - Surgery, which is potentially risky
  - Medical management, which has a low success rate
- With surgery, one must assess the extent of disease and decide between curative and palliative surgery
- ***Goal: maximize life expectancy for the patient***

The initial decision is between surgery and medical management



Treatment is initiated on patients w/ symptoms; some w/o disease



Those with disease have a chance to benefit from treatment



Likewise with surgery



Surgery is risky even for those with no disease




For disease, try cure vs. palliate?
Try
Cure
Palliate
Disease
Present
Surgery
Live
Disease
Absent
Surg
Dth
Cure
Disease
Present
No
Cure
Medical
Mgmt
Disease
Absent


Surgical risks here too
Surgery
Disease
Present
Disease
Absent
Live
Surg
Dth
Try
Cure
Surg
Dth
Live
Palliate
Surg
Dth
Live
Medical
Mgmt
Disease
Present
Cure
No
Cure
Disease
Absent

Chance of cure for survivors




Paths define course of events
Surgery
Disease
Present
Disease
Absent
Live
Surg
Dth
Cure
Try
Cure
Surg
Dth
Cure
Live
No
Cure
Surg
Dth
Palliate
Cure
Medical
Mgmt
Disease
Present
Disease
Absent
1. Surgery
2. in a patient with disease
3. where curative surgery chosen
4. and patient survives
5. and is cured


Surgery
Disease
Present
10%
90%
Disease
Absent
Live
99%
1%
Surg
Dth
Try
Cure
Surg
Dth
10%
90%
Live
Cure
90%
10%
No
Cure
Palliate
Surg
Dth
2%
98%
Live
Cure
10%
90%
No
Cure
Medical
Mgmt
Disease
Present
10%
90%
Disease
Absent
Cure
10%
90%
No
Cure
Add probabilities
(studies, experts)


Surgery
Disease
Present
10%
90%
Disease
Absent
Live
99%
1%
Surg
Dth
Try
Cure
Surg
Dth
10%
90%
Live
Cure
90%
10%
No
Cure
Palliate
Surg
Dth
2%
98%
Live
Cure
10%
90%
No
Cure
Medical
Mgmt
Disease
Present
10%
90%
Disease
Absent
Cure
10%
90%
No
Cure
Now add outcomes



Death yields 0 years of additional life


Surgery
Disease
Present
10%
90%
Disease
Absent
Live
99%
1%
Surg
Dth
0 Y
Medical
Mgmt
Disease
Present
10%
Cure
10%
90%
No
Cure
2 Y
90%
Disease
Absent
Try
Cure
Surg
Dth
10%
0 Y
Cure
90%
90%
Live
10%
No
Cure
2
Palliate
Surg
Dth
2%
0 Y
98%
Live
Cure
10%
90%
No
Cure
2
Uncured disease confers 2 years of additional life


Surgery
Disease
Present
10%
90%
Disease
Absent
Live
99%
20 Y
1%
Surg
Dth
0 Y
Try
Cure
Surg
Dth
10%
0 Y
Cure
90%
20
90%
Live
10%
No
Cure
2
Palliate
Surg
Dth
2%
0 Y
98%
Live
Cure
10%
20
90%
No
Cure
2
Medical
Mgmt
Disease
Present
10%
Cure
10%
20 Y
90%
No
Cure
2 Y
90%
Disease
Absent
20 Y
Cure yields 20 years
of additional life

Now average out & fold back



Now average out & fold back



10%*20 + 90%*2 =
3.8 years (expected)

Now average out & fold back



Same calculation here

Now average out & fold back



10%*20 + 90%*2 =
3.8 years (expected)

Now average out & fold back



Since disease presence unknown, we do this again

Now average out & fold back



10%*3.8 + 90%*20 =
18.38 years

Now average out & fold back



Now average out & fold back



Now average out & fold back



Now average out & fold back



Now average out & fold back



Now average out & fold back





This one is different:
Decision node:
Surgeon picks option with greatest expected benefit:
Try Cure (16.38 years) preferred
(called "folding back")

Now average out & fold back



Now average out & fold back



Now average out & fold back




Surgery
19.46 Y
Medical
Mgmt
18.38 Y



Decision node again (overall)
Surgery is preferred to Medical Management because the incremental benefit of surgery is:

19.46 – 18.38 = **1.08 years**

Recommendation: Choose surgery (with "try cure" surgical option)



Use same approach for CEA but now with second set of outcomes

19.46 – 18.38 = **1.08 years**

$10,000 – $100 = **$9,900**

$9,900 / 1.08 =

$9,167 per life year gained

Surgery if willing to pay at least $9,167 per life year gained, otherwise medical management

SENSITIVITY ANALYSIS


Surgery
Disease
Present
10%
90%
Disease
Absent
Live
99%
20 Y
1%
Surg
Dth
0 Y
Try
Cure
Surg
Dth
10%
0 Y
Cure
90%
20
Live
90%
10%
No
Cure
2
Palliate
Surg
Dth
2%
0 Y
Cure
98%
10%
20
Live
90%
No
Cure
2
Medical
Mgmt
Disease
Present
10%
Cure
10%
20 Y
90%
No
Cure
2 Y
90%
Disease
Absent
20 Y
But probabilities and
Outcome values
uncertain…

Sensitivity Analysis

- Systematically asking “what if” questions to see how the decision result changes
- Determines how “robust” the decision is
  - **Threshold analysis:** one parameter varied
  - **Multi-way analysis:** multiple parameters systematically varied


Surgery
Disease
Present
10%
90%
Disease
Absent
Live
99%
20 Y
1%
Surg
Dth
0 Y
Medical
Mgmt
Disease
Present
10%
Cure
10%
20 Y
90%
No
Cure
2 Y
90%
Disease
Absent
20 Y
Try
Cure
Surg
Dth
10%
0 Y
Cure
90%
20
Live
90%
10%
No
Cure
2
Palliate
Surg
Dth
2%
0 Y
98%
Live
Cure
10%
20
90%
No
Cure
2
If probability of surgical death with curative surgery uncertain


1-way sensitivity analysis: Curative Surgical Death
Base Case
Threshold
20.000
19.500
19.000
18.500
18.000
17.500
Expected Life Years
0.00
0.10
0.20
0.30
0.40
0.50
0.60
0.70
0.80
0.90
1.00
Probability of Curative Surgical Death
Surgery - Curative
Med Mgmt


Medical Management Preferred
Surgery Preferred
Base Case
1.00
0.90
0.80
0.70
0.60
0.50
0.40
0.30
0.20
0.10
0.00
Probability of Curative Surgical Death
0.00
0.10
0.20
0.30
0.40
0.50
0.60
0.70
0.80
0.90
1.00
Prevalence of Disease

POLL SLIDE

- **Sensitivity analyses tell us (*choose all answers that you believe to be correct*):**
  1. How much our model outputs change based on changes in our model inputs
  2. Whether or not we would change our decision if our model inputs were different
  3. How uncertain we are about making the decision based on the model
  4. Tell us whether the decision-problem is politically or ethically sensitive

Advanced: Probabilistic Sensitivity Analysis (2nd order Monte Carlo)

- Decision tree estimates of probabilities and utilities are replaced with probability distributions (e.g. logistic-normal)
- The tree is evaluated *many* times with random values selected from each distribution
- Results include means and standard deviations of the expected values of each strategy

MARKOV MODELS VS. DECISION TREES

WHAT TO DO WHEN THERE IS A POSSIBILITY OF REPEATED EVENTS AND/OR DECISIONS?

Decision about one-time, immediate action



Decision about one-time, immediate action



Decisions: repeated actions and/or with time-dependent events



Repeated in what sense?



Disease process involves events occurring at multiple time points



Intervention (can) be delivered repeatedly too



- Repeated events can occur throughout an individual's life.
- Interventions delivered at multiple time points. Subsequent transitions depend on prior intervention outcomes.

What is a Markov Model?

- **Markov Model:** Mathematical modeling technique, derived from matrix algebra, that describes the transitions a cohort of patients make among a number of mutually exclusive and exhaustive health states during a series of short intervals or cycles

Properties of a Markov Model

- Individuals are always in one of a finite number of health states
- Events are modeled as transitions from one state to another
- Time spent in each health state determines overall expected outcome
  - Living longer without disease yields higher life expectancy and quality adjusted life expectancy
- During each cycle of the model, individuals may make a transition from one state to another

Constructing a Markov Model

- Define mutually exclusive health states
- Determine possible transitions between these health states
  - State transitions
  - Transition probabilities
- Determine clinically valid cycle length

Cycle Length

- Short enough that for a given disease being modeled the chance of two events/transitions occurring in one cycle is essentially 0
  - Many applications: weekly or monthly
  - Some (e.g., ICU) may hourly or daily

Natural history disease model: health states

- Mutually exclusive and collectively exhaustive health states
- Best defined by actual biology/pathophysiology
- Markovian assumptions:
  - Homogeneity: All individuals in the same state have the same costs, quality of life, risks of transition
  - Memorilessness: The current state determines future risks
  - **Note: Stratification and tunnel states used to ensure Markov assumptions hold (advanced topic)**

Natural history disease model: transitions



- Transitions between health states (arrows)
- The proportion that do not transition stay in current state
- Risk of death at all times and from all states!
- If no transition out of a state = absorbing state (i.e., death)

Natural history disease model: time and matrix representation



$$\begin{pmatrix} pHH & pSH & 0 \\ pHS & pSS & 0 \\ pHD & pSD & 1 \end{pmatrix}$$

For example pSH is the Probability of going from Sick to Healthy

Natural history disease model: time and matrix representation



$$\begin{pmatrix} pHH & pSH & 0 \\ pHS & pSS & 0 \\ pHD & pSD & 1 \end{pmatrix} \begin{pmatrix} propH \\ propS \\ propD \end{pmatrix}$$

time=t

At time t, cohort has proportions in various states (Sum to 1!)

Natural history disease model: time and matrix representation



$$\begin{pmatrix} pHH & pSH & 0 \\ pHS & pSS & 0 \\ pHD & pSD & 1 \end{pmatrix} \begin{pmatrix} propH \\ propS \\ propD \end{pmatrix}_{time=t} = \begin{pmatrix} propH \\ propS \\ propD \end{pmatrix}_{time=t+1}$$

NOTE: transition probabilities can be time dependent as well

Natural history disease model: time and matrix representation



$$\begin{pmatrix} pHH & pSH & 0 \\ pHS & pSS & 0 \\ pHD & pSD & 1 \end{pmatrix} \begin{pmatrix} propH \\ propS \\ propD \end{pmatrix}_{time=t} = \begin{pmatrix} propH \\ propS \\ propD \end{pmatrix}_{time=t+1}$$

Natural history disease model: time and matrix representation



$$\begin{pmatrix} pHH & pSH & 0 \\ pHS & pSS & 0 \\ pHD & pSD & 1 \end{pmatrix} \begin{pmatrix} propH \\ propS \\ propD \end{pmatrix}_{time=t} = \begin{pmatrix} propH \\ propS \\ propD \end{pmatrix}_{time=t+1}$$

Model trace



Model trace



- Is proportion the prevalence?
- Is model time the age?

Underlying the trace

Stage	propH_t	propS_t	propD_t	NotD
0	1.00	0.00	0.00	1.00
1	0.90	0.09	0.01	0.99
2	0.75	0.10	0.15	0.85
3	0.50	0.25	0.25	0.75
4	0.20	0.40	0.40	0.60
5	0.10	0.30	0.60	0.40
6	0.05	0.15	0.80	0.20
7	0.00	0.00	1.00	0.00

Quality Adjusted Life Years (QALYS) & quality-of-life weights



Valuing outcomes



Stage	propH_t	propS_t	propD_t	NotD
0	1.00	0.00	0.00	1.00
1	0.90	0.09	0.01	0.99
2	0.75	0.10	0.15	0.85
3	0.50	0.25	0.25	0.75
4	0.20	0.40	0.40	0.60
5	0.10	0.30	0.60	0.40
6	0.05	0.15	0.80	0.20
7	0.00	0.00	1.00	0.00

$$QALYs = \sum_{t=0}^{T} \left[\left(propH_t * qH \right) + \left(propS_t * qS \right) + \left(propD_t * 0 \right) \right]$$

$$COSTs = \sum_{t=0}^{T} \left[\left(propH_t * cH \right) + \left(propS_t * cS \right) + \left(propD_t * 0 \right) \right]$$

Interventions?



$$\begin{pmatrix} pHH & pSH & 0 \\ pHS & pSS & 0 \\ pHD & pSD & 1 \end{pmatrix} \begin{pmatrix} propH \\ propS \\ propD \end{pmatrix}_{time=t} = \begin{pmatrix} propH \\ propS \\ propD \end{pmatrix}_{time=t+1}$$

Screening before treatment

- Screening 70% sensitivity, 100% specific
- Treatment 90% effective
- Intervention occurs after natural hx transitions every cycle

- Calculations
  - pHS_i = pHS*(0.3) + pHS*(0.7*0.1)
  - pSS_i = pSS*(0.3) + pSS*(0.7*0.1)
  - pSH_i = pSH + pSS*(0.7*0.9)
  - pHH_i = pHH + pHS*(0.7*0.9)

Natural History

$$\begin{pmatrix} 0.5 & 0.2 & 0 \\ 0.4 & 0.6 & 0 \\ 0.1 & 0.2 & 1 \end{pmatrix}$$

Screening before treatment

pHH_i	pSH_i	0
pHS_i	pSS_i	0
pHD	pSD	1

Screening before treatment

0.752	0.222	0
0.148	0.578	0
0.100	0.200	1

With and w/o intervention



With and w/o intervention



Natural History



Intervention



Intervention



Cohorts vs. individuals
Deterministic vs. stochastic

- Markov cohort model (i.e., the matrix version) is smooth model (infinite population size) of the proportion of a cohort in each state at each time
- Can use same structure to simulate many individuals (first-order Monte Carlo) (simple microsimulation)
- The matrix becomes the probability of an individual transition from one state to another instead of the % of those in a given state who deterministically flow into another state

Microsimulation



Microsimulation



Microsimulation



Microsimulation



Recall the trace and calculation of outcomes from it

Stage	propH_t	propS_t	propD_t	NotD
0	1.00	0.00	0.00	1.00
1	0.90	0.09	0.01	0.99
2	0.75	0.10	0.15	0.85
3	0.50	0.25	0.25	0.75
4	0.20	0.40	0.40	0.60
5	0.10	0.30	0.60	0.40
6	0.05	0.15	0.80	0.20
7	0.00	0.00	1.00	0.00

$$QALYs = \sum_{t=0}^{T} \left[\left(propH_t * qH \right) + \left(propS_t * qS \right) + \left(propD_t * 0 \right) \right]$$

$$COSTs = \sum_{t=0}^{T} \left[\left(propH_t * cH \right) + \left(propS_t * cS \right) + \left(propD_t * 0 \right) \right]$$

Microsimulation

- Run with many individuals
- Calculate proportions in each state at each time (just like in our Markov cohort table)
  - Stage 2: 5100 sick / 100,000 people = 5.1%
- Approximates the “smooth” cohort version
  - 5.1% [CI] is ~= 5.0% in “smooth” cohort
  - **<u>Advanced</u>**
    - Larger the number of individuals the closer to the smooth cohort (tighter the CI)
    - See Kuntz/Weinstein chapter of Michael Drummond’s book on Economic Evaluation for more on this for more on this

Why consider microsimulation?

- It requires longer simulation times
- It is more complex
- Fewer people are familiar with it
- There is “Monte Carlo” noise (random error) even with simulating fairly large groups of individuals (at least for rare events)

State explosion!

- Suppose you want to use a Markov model of a disease with 2 states and death (H,S,D)
- Suppose you need it stratified by sex and smoking status (3 levels), BMI (4 levels), hypertension (4 levels)
- Now you need 2x3x4x4x2 states (death is not stratified = 192 states
- What if you need to stratify states by past history? (previous high hypertension, used to be obese) or Tx history (has a stent)?

Microsimulation as alternative

- Simulate 1 individual at a time
- Assign a set of attributes to the individual
  - Sex=M, Smoking=Y, BMI=Overweight, HT=Y
- Define a function for the probability of transitioning from H to S
  - P(H to S | Sex, Smoking, BMI, HT)
- Have functions for changing attributes
  - P(BMI=Obese|Sex, BMI)
- Track previous health states
  - P(H to S| Sex, Smoking, BMI, HT, S in the past)
- **Note: Could estimate these functions from logistic regressions**

Sage advice I have heard

- Know what information your consumers need
- Pick a model that is as simple as possible … but no simpler
- Know the limits of what your model does and make statements within those limits – All research studies have limitations

Summary:
Medical Decision Analysis

- Clearly defines alternatives, events, and outcomes
- Formal method to combine evidence
- Can prioritize information acquisition
- Can help healthcare providers to make medical decisions under uncertainty

Classic sources on about decision analysis and modeling

- Sox HC, Blatt MA, Higgins MC, Marton KI (1988) Medical Decision Making. Boston MA: Butterworth-Heinemann Publisher.
- Detsky AS, Naglie G, Krahn MD, Naimark D, Redelmeier DA. Primer on medical decision analysis: Parts 1-5. Med Decis Making. 1997;17(2):123-159.
- Sonnenberg FA, Beck JR. Markov models in medical decision making: a practical guide. Med Decis Making. 1993;13(4):322-38.
- Beck JR, Pauker SG. The Markov process in medical prognosis. Med Decis Making. 1983;3(4):419-458.
- Society for Medical Decision Making (http://www.smdm.org)

THANK YOU

Jeremy Goldhaber-Fiebert
(jeremygf@stanford.edu)

HEALTH POLICY STANFORD

CENTER FOR HEALTH POLICY/
CENTER FOR PRIMARY CARE AND OUTCOMES RESEARCH